

Wiedemann-Franz Law - Answer Sheet

Part A: Electric conductivity of copper, aluminum and brass (1.5 points)

A.1 (1.0 pt)

Magnet descend time:

number	Copper	Aluminum	Brass


A.2 (0.5 pt)

	Copper	Aluminum	Brass
Electrical conductivity			

Part B: Thermal conductivity of copper (3.0 points)

B.1 (0.1 pt)


Rod 1 temperature :

**B.2** (0.5 pt)**B.3** (0.1 pt) $P =$ **B.4** (0.5 pt)

Time	T_1	T_2	T_3	T_4	T_5	T_6	T_7	T_8

B.5 (1.0 pt)

Draw in the additional graph papers the temperature as a function of location.



B.6 (0.5 pt)

$$\kappa_0 =$$

$$\frac{\Delta T}{\Delta t} =$$

B.7 (0.3 pt)


Circle the correct answer:

$$\kappa > \kappa_0 \text{ or } \kappa < \kappa_0 \text{ or } \kappa = \kappa_0$$

Part C: Estimating the heat loss and the heat capacity of copper (4.0 points)

C.1 (1.0 pt)

[illegible]

**C.2** (1.0 pt)

Draw in the additional graph papers the average temperature as function of time

C.3 (1.0 pt)

Expression:

$$c_p =$$

$$\dot{Q}_{loss} =$$

Value:

$$c_p =$$

$$\dot{Q}_{loss} =$$

C.4 (1.0 pt)

Expression:

$$\kappa_{copper} =$$


Value:

$$\kappa_{copper} =$$

Part D: Measure the heat conductivity of brass and aluminum (1.0 points)**D.1** (0.1 pt)

Rod 2 : $T =$

Experiment



A2-6

English (Official)

D.2 (0.2 pt)

Reading time :

T_1	T_2	T_3	T_4	T_5	T_6	T_7	T_8

$\Delta T_{Copper-1} / \Delta x$	$\Delta T_{Brass} / \Delta x$	$\Delta T_{Aluminum} / \Delta x$	$\Delta T_{Copper-2} / \Delta x$

D.3 (0.7 pt)

Expression:

$$\kappa_{Aluminum} =$$

$$\kappa_{Brass} =$$

Value:

$$\kappa_{Aluminum} =$$

$$\kappa_{Brass} =$$

Part E: Wiedemann-Franz law (0.5 points)

E.1 (0.5 pt)

	Copper	Aluminum	Brass
Electrical conductivity			
Heat conductivity			
Lorenz coefficient			

